

Amara fallax, extremely common in garden. This insect is on the wing a great deal on hot days, when it may be picked up anywhere, just after alighting.

Bembidion erosum, Bembidion zephyrum, from sandy beaches.

Bembidion dubitans, garden, not frequent.

Bembidion nigripes, garden, very common, both species taken running over open ground.

Carabus nemoralis. I have some evidence that this widespread species may be patchy in distribution. Though I have never taken it here, it appears to be fairly common at Nanaimo only seven miles distant. Among dead leaves and moss on ground.

Carabus lapelayi. Two specimens taken from under boards in garden during June, Wellington.

Dicheirus piceus. Very common on Rocky barren shorelines, under stones and logs. An attempt at collecting Coleoptera on several dry rocky islands in the Strait of Georgia, resulted in this species only being taken in any numbers. In addition, I had only one Harpalus cautus, and one large Staphylinidae, (Ocypus ater?) Batches of D. piceus taken in late September contained many teneral individuals.

Zacotus matthewsi. One only from sandy beach, under log, Tofino, May 16th. Collected by Kenneth Gibson.

Bothriopterus adstrictus. Very common under boards in garden, not taken elsewhere.

OBITUARY

In Dr. Daniel Louis Uyttenboogaart, who passed away after a painful illness on August 2nd, 1947 in his seventy-fifth year, the Nederlandsche Entomologische Vereeniging (Dutch Entom. Soc.) has lost its President and one of the most prominent Coleopterists among its members. He was a well known specialist in Curculionidae and in Coleoptera from the Azores. He made collecting trips to the Azores and to South Africa.

P. H. v. Doesburg.
Baarn - HOLLAND
